

GUJARAT TECHNOLOGICAL UNIVERSITY

BE - SEMESTER-I & II (NEW) EXAMINATION – SUMMER-2019

Subject Code: 2110015

Date: 01/06/2019

Subject Name: Vector Calculus & Linear Algebra

Time: 10:30 AM TO 01:30 PM

Total Marks: 70

Instructions:

1. Question No.1 is compulsory. Attempt any four out of remaining six questions.
2. Make suitable assumptions wherever necessary.
3. Figures to the right indicate full marks.

Q.1 Objective Question (MCQ)

(a)

07

1. The matrix $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$ is in the form
(a) Row echelon. (b) Reduced row echelon. (c) Both (a) and (b). (d) None.
2. For $A = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$ the $|A^k| =$ _____
(a) 1 (b) 2 (c) 2^k (d) 2^{k-1}
3. If u and v are vectors in a real inner product space, and $\|u\|=2$, $\|v\|=3$, then $|\langle u, v \rangle| \leq$ _____
(a) 6 (b) 3 (c) 2 (d) 1.5
4. Which of the following doesn't lie in the space spanned by $\cos^2 x$ and $\sin^2 x$?
(a) 1 (b) 0 (c) $\sin x$ (d) $\cos 2x$
5. Dimension of the subspace $\{ p(x) \in P_2 : p(0) = 0 \}$ of $P_2 = \{ a+bx+cx^2 : a, b, c \in \mathbb{R} \}$ is
(a) 3 (b) 2 (c) 1 (d) 0
6. Which of the following subsets of $\mathbb{R}^2$ is linearly dependent?
(a) $\{(1,2), (2,1)\}$ (b) $\{(1,2), (2,1), (1,1)\}$ (c) $\{(1,2)\}$ (d) None
7. Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(x,y) = (x,0)$ then $\text{Ker}(T) =$ _____
(a) Y-axis (b) X-axis (c) Origin (d) None

(b)

07

1. Which of the following is not an elementary matrix?
(a) $\begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$ (b) $\begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$ (c) $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 9 \\ 0 & 0 & 1 \end{pmatrix}$ (d) $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$
2. For $\vec{a} = (1, -1, 2)$, $\vec{b} = (1, 3, 1)$ are vectors of $\mathbb{R}^3$ with Euclidean inner product then $\cos \theta =$ _____, where θ is the angle between the two vectors.
(a) 1 (b) 0 (c) -3 (d) 6
3. Which of the following is not true?
(a) $(AB)^T = B^T A^T$ (b) $(AB)^{-1} = B^{-1} A^{-1}$ (c) $(A^T)^T = A$ (d) $A^T = -A$
4. If A is $n \times n$ matrix having rank $n-1$ then $A, A^2, A^3, \dots, A^k, \dots$ have common eigenvalue _____
(a) 1 (b) -1 (c) 0 (d) 2
5. If A is unitary matrix then $A^{-1} =$ _____
(a) A (b) A^2 (c) A^T (d) I
6. The dimension of the solution space of $x - y = 0$ is _____
(a) 0 (b) 1 (c) 2 (d) 3
7. If $f(x,y,z) = xyz$ then $\text{Curl}(\text{grad } f) =$ _____
(a) 0 (b) x (c) $xi+yj+zk$ (d) xyz

- Q.2 (a)** Which of the following are linear combination of $u = (0, -2, 2)$ and $v = (1, 3, -1)$? Justify! (i) $(2, 2, 2)$, (ii) $(0, 4, 5)$ **03**
- (b)** Using Gram-Schmidt orthogonalization process find the corresponding orthonormal set to $\{(1, 1, 1), (0, 1, 1), (0, 0, 1)\}$. **04**
- (c)** Using Gauss- Jordan elimination find the inverse of $\begin{pmatrix} -1 & 3 & -4 \\ 2 & 4 & 1 \\ -4 & 2 & -8 \end{pmatrix}$. **07**
- Q.3 (a)** Find the rank of the matrix and basis of the null space of $\begin{pmatrix} 1 & -1 & 3 \\ 5 & -4 & -4 \\ 7 & -6 & 2 \end{pmatrix}$. **03**
- (b)** Solve the system of linear equations using Gauss elimination method: $x + y + 2z = 8, -x - 2y + 3z = 1, 3x - 7y + 4z = 10$. **04**
- (c)** Show that the set of all real numbers of the form $(x, 1)$ with operations $(x, 1) + (x', 1) = (x + x', 1)$ and $k(x, 1) = (kx, 1)$ forms a vector space. **07**
- Q.4 (a)** Determine whether the following are linear transformation or not? **03**
- (i) $T: P_2 \rightarrow P_2, T(p(x)) = p(x + 1)$,
- (ii) $T: P_2 \rightarrow P_2, T(a + bx + cx^2) = (a + 1) + (b + 1)x + (c + 1)x^2$.
- (b)** Which of the following sets of vectors of R^3 are linearly independent? Justify. **04**
- (i) $\{(4, -1, 2), (-4, 10, 2)\}$ (ii) $\{(-3, 0, 4), (5, -1, 2), (1, 1, 3)\}$
- (c)** Find the eigenvalues and bases for the eigenspaces for A^{11} , **07**
- $$A = \begin{pmatrix} -1 & -2 & -2 \\ 1 & 2 & 1 \\ -1 & -1 & 0 \end{pmatrix}$$
- Q.5 (a)** Find basis of kernel and range of $T: R^2 \rightarrow R^2$, defined by $T(x, y) = (2x - y, -8x + 4y)$ **03**
- (b)** Which of the following are basis of R^3 ? Justify! **04**
- (i) $\{(1, 0, 0), (2, 2, 0), (3, 3, 3)\}$, (ii) $\{(3, 1, -4), (2, 5, 6), (1, 4, 8)\}$
- (c)** Let $T: P_2 \rightarrow P_2$, defined by $T(p(x)) = p(3x - 5)$ **07**
- (i) Find the matrix of T with respect to the basis $\{1, x, x^2\}$.
- (ii) Use the indirect procedure using matrix to compute $T(1 + 2x + 3x^2)$.
- (iii) Check the result in (b) by computing $T(1 + 2x + 3x^2)$ directly.
- Q.6 (a)** Show that $\vec{F} = \frac{(y\vec{i} - x\vec{j})}{x^2 + y^2}$ is irrotational. **03**
- (b)** Find the directional derivative of $f(x, y, z) = x^2z + y^3z^2 - xyz$ at $(1, 1, 1)$ in the direction of the vector $(-1, 0, 3)$. **04**
- (c)** Using Green's theorem evaluate $\oint_C (3x^2 - 8y^2) dx + (4y - 6xy) dy$, where C is the boundary of the region bounded by $y^2 = x$ and $y = x^2$. **07**
- Q.7 (a)** Find the work done by $\vec{F} = (y - x^2)\vec{i} + (z - y^2)\vec{j} + (x - z^2)\vec{k}$ over the curve $\vec{r}(t) = t\vec{i} + t^2\vec{j} + t^3\vec{k}$; $0 \leq t \leq 1$, from $(0, 0, 0)$ to $(1, 1, 1)$. **03**
- (b)** Use Cramer's rule to solve: $x + 2z = 6, -x + 4y + 6z = 30, -x - 2y + 3z = 8$. **04**
- (c)** Verify divergence theorem for $\vec{F} = x\vec{i} + y\vec{j} + z\vec{k}$ over the sphere $x^2 + y^2 + z^2 = a^2$. **07**